

Mandelbrot Set

Keywords: complex numbers, Gaussian plane, sequences, convergence, absolute value

The Mandelbrot set is one of the most famous and beautiful fractals, fascinating mathematicians, scientists, and artists around the world. Although it looks like a complex pattern at first glance, it is based on a simple mathematical rule of repeated exponentiation and addition. What makes it so interesting is its infinite complexity and the beautiful patterns hidden in every detail.




Figure 1: The Mandelbrot set; the color of points in its vicinity corresponds to the index of the first term in the sequence at which it is determined that the sequence goes to infinity.

The use of the Mandelbrot set goes far beyond mathematics. It is used in computer graphics to create complex and realistic objects when modeling natural structures such as coastlines, mountains, or cloud formations. It can even be used in economics and physics to help simulate chaotic systems.

The Mandelbrot set is proof that even simple mathematical procedures can lead to incredibly complex and beautiful results with real-world applications.

Creating a Set

Let us consider a relatively simple recurrent formula

$$z_{n+1} = z_n^2 + c,$$

where the initial value is $z_0 = 0$ and c represents an arbitrary complex number. The French-American mathematician Benoit Mandelbrot (1924–2010) was interested in when the sequence of numbers formed this way is bounded, i.e. for which c in the complex plane the sequence converges or oscillates. If at some point the sequence diverges, he wanted to know how fast. It can be proven that if the absolute value of any term of the sequence z_n exceeds 2, then this sequence is not bounded.

The Mandelbrot set is then the set of points c in the complex plane for which the sequence created by the recurrent formula converges or oscillates. Thanks to the above fact, we know that for each term z of this sequence, its absolute value $|z|$ must be less than or equal to two.

Verification of whether a given c belongs to the Mandelbrot set is carried out by calculating individual iterations and observing the absolute values of these iterations. To calculate the iterations, we use the recurrent formula

$$z_{n+1} = z_n^2 + c, \quad z_0 = 0.$$

For example, for $c = -i$ we obtain:

$$\begin{aligned} z_1 &= z_0^2 - i = 0^2 - i = -i, & |z_1| &= 1, \\ z_2 &= z_1^2 - i = (-i)^2 - i = -1 - i, & |z_2| &= \sqrt{2}, \\ z_3 &= z_2^2 - i = (-1 - i)^2 - i = i, & |z_3| &= 1, \\ z_4 &= z_3^2 - i = (i)^2 - i = -1 - i, & |z_4| &= \sqrt{2}. \end{aligned}$$

It is clear from the calculation that the results $-i$ and $-1 - i$ will repeat indefinitely. Therefore, the condition $|z| \leq 2$ will always be satisfied, and therefore the number $-i$ belongs to the Mandelbrot set.

Exercises

Exercise 1. Verify whether the complex numbers 1 ; i ; -1 ; $1 + i$ belong to the Mandelbrot set.

Exercise 2. Prove that if there exists $k \in \mathbb{N}$ such that $|z_k| > 2$, then the sequence z_n diverges.

Literature

- Čápka Hartinger, David. *Mandelbrotova množina - lekce 3 [online]* <https://www.itnetwork.cz/algoritmy/graficke/algoritmy-graficke/mandelbrotova-mnozina> [cit. 22. 9. 2023]
- Wikipedia. *Mandelbrot set [online]*. Available from https://en.wikipedia.org/wiki/Mandelbrot_set [cit. 22. 9. 2023].
- PantheraLeo1359531. *Mandelbrot set - figure [online]*. Available from https://commons.wikimedia.org/w/index.php?title=Mandelbrot_set&file=Mandelbrot_set.jpg [cit. 22. 9. 2023]